

LABELLING THE CHARACTER TABLES OF SYMMETRIC AND ALTERNATING GROUPS

MARK WILDON

ABSTRACT. Let X be a character table of the symmetric group S_n . It is shown that unless $n = 4$ or $n = 6$, there is a unique way to assign partitions of n to the rows and columns of X so that for all λ and ν , $X_{\lambda\nu}$ is equal to $\chi^\lambda(\nu)$, the value of the irreducible character of S_n labelled by λ on elements of cycle type ν . Analogous results are proved for alternating groups, and for the Brauer character tables of symmetric and alternating groups.

1. INTRODUCTION

In [10] Nagao proved that if G is a finite group with a character table which differs from a character table of the symmetric group S_n only by a permutation of its rows and columns, then G is isomorphic to S_n . In this paper we consider a question naturally raised by Nagao's result. To state it, we must recall that the ordinary irreducible characters of S_n are canonically labelled by the partitions of n , and that this set also labels the conjugacy classes of S_n . Given partitions λ and ν of n , let $\chi^\lambda(\nu)$ be the value of the irreducible character of S_n labelled by λ on elements of cycle type ν .

Now suppose that one has discovered (for example, by applying Nagao's theorem) that a given square matrix is an unlabelled character table of the symmetric group S_n . We ask: when can one go further, and *uniquely* reconstruct the partitions labelling its rows and columns? The answer is given by the following theorem.

Theorem 1.1. *Let X be a character table of the symmetric group S_n . Unless $n = 4$ or $n = 6$, there is a unique way to assign partitions of n to the rows and columns of X so that $X_{\lambda\nu} = \chi^\lambda(\nu)$ for all partitions λ, ν of n . If $n = 4$ or $n = 6$ then there are exactly two different labellings.*

Probably the reader already knows the reason for one of the exceptional cases. We discuss them in §2.1. The main work begins in §2.2, where we show that, provided $n \geq 7$, there is only one possible way to assign partitions of the form $(n - m, m)$ to the rows of X . (We say that such partitions are *two-row partitions*.) Then in §2.3 we show that given such a partial row labelling, there is only one way to assign all the column labels. Of course, once we have fixed the column labels, the remaining row labels are uniquely determined. It follows that there is a unique way to label X . In §2.4 we give an efficient way to complete the row labelling.

Date: October 12, 2006

2000 Mathematics Subject Classification 20C30 (primary), 20C20 (secondary).

It is natural to ask the analogous question for alternating groups, and for the Brauer character tables of symmetric and alternating groups. Strikingly, these questions may also be answered using the strategy we have just outlined, giving the results stated below in Theorem 3.1, Theorem 4.1 and Theorem 5.2 respectively. (We only deal with alternating groups in odd characteristic.)

Informally, we may interpret our results as saying that the rich structure of representations of symmetric and alternating groups gives their character tables a high degree of rigidity. In §6 we make this idea more precise by putting our results in the general context of representations of finite groups. We also pose two open problems suggested by our work.

2. PROOF OF THEOREM 1.1

2.1. We begin with S_6 , the only symmetric group to have an outer automorphism (see for example [12, Theorem 7.9]). Any outer automorphism of S_6 permutes the conjugacy classes of S_6 by

$$(6) \leftrightarrow (3, 2), (3, 3) \leftrightarrow (3), (2, 2, 2) \leftrightarrow (2)$$

and permutes its ordinary characters by

$$(5, 1) \leftrightarrow (2, 2, 2), (2, 1, 1, 1, 1) \leftrightarrow (3, 3), (4, 1, 1) \leftrightarrow (3, 1, 1, 1).$$

Thus if σ and τ denote the corresponding permutations on the set of partitions of 6 then $\chi^\lambda(\nu) = \chi^{\lambda^\sigma}(\nu^\tau)$ for all partitions λ, ν . This gives two different ways to label the character table of S_6 . Inspection of the table shows there are no more.

We now turn to S_4 , which has the character table shown below.

	(1^4)	$(2, 1, 1)$	(4)	$(2, 2)$	$(3, 1)$
(4)	1	1	1	1	1
$(3, 1)$	3	-1	1	-1	0
$(2, 1, 1)$	3	1	-1	-1	0
$(2, 2)$	2	0	0	2	-1
(1^5)	1	-1	1	1	-1

Notice that if we swap the columns labelled by $(2, 1, 1)$ and (4) , and the rows labelled by $(3, 1)$ and $(2, 1, 1)$, we end up with the same matrix. Again it is easy to see that this is the only alternating way to label the character table. Unlike the case of S_6 , this alternative labelling is not induced by any automorphism of the group — its existence appears to be entirely coincidental. (See §6 for some remarks related to this phenomenon.)

Theorem 1.1 may readily be verified by inspection of the character tables if $n \leq 3$ or $n = 5$, so from now on we shall assume $n \geq 7$.

2.2. Let $n \geq 7$ and let X be an unlabelled character table of S_n . We shall show that there is a unique way to assign two-row partitions (that is, partitions of the form $(n - m, m)$) to the rows of X .

By the orthogonality relations for ordinary characters, there is a unique row of X containing only positive entries. Similarly there is a unique column of X containing only positive entries. We may therefore uniquely identify the column of the character table corresponding to the identity element and

the row corresponding to the trivial character, thus fixing the row label (n) and the column label (1^n) .

For all $n \neq 6$ the symmetric group S_n has exactly two characters of degree $n - 1$, namely $\chi^{(n-1,1)}$ and $\chi^{(2,1^{n-2})}$ (see [6] Theorem 2.4.10). These characters are defined by

$$\chi^{(n-1,1)}(g) = |\text{Fix } g| - 1,$$

where $\text{Fix } g$ is the set of elements fixed by g in its natural action on $\{1, \dots, n\}$, and

$$\chi^{(2,1^{n-2})}(g) = \text{sgn}(g)\chi^{(n-1,1)}(g).$$

For $n > 4$, only $\chi^{(n-1,1)}$ takes $n - 3$ as a value. Hence we may fix the row label $(n - 1, 1)$.

To proceed further, we observe that given two rows in the unlabelled character table X we may multiply the corresponding entries and so obtain a new character. Then, still working only with the unlabelled table, we may take the inner product of this character with each row, and so determine its irreducible constituents. The following lemma gives the results needed to exploit these observations. (The main idea in the proof of this lemma comes from [1].)

Lemma 2.1. *If $n \geq 4$ then*

$$\chi^{(n-1,1)}\chi^{(n-1,1)} = \chi^{(n)} + \chi^{(n-1,1)} + \chi^{(n-2,2)} + \chi^{(n-2,1,1)}.$$

If $r \geq 2$ and $n > 2r + 1$ then

$$\begin{aligned} \chi^{(n-1,1)}\chi^{(n-r,r)} &= \chi^{(n-r-1,r+1)} + \chi^{(n-r,r)} \\ &\quad + \chi^{(n-r+1,r-1)} + \chi^{(n-r-1,r,1)} + \chi^{(n-r,r-1,1)}. \end{aligned}$$

Proof. Recall that $\chi^{(n-1,1)} + \chi^{(n)}$ is the permutation character of S_n acting on $\{1, \dots, n\}$. As this character is induced from the subgroup S_{n-1} of S_n , we have

$$\chi^{(n-1,1)}\theta = \left(1_{S_{n-1}} \uparrow^{S_n}\right)\theta - \theta = \theta \downarrow_{S_{n-1}} \uparrow^{S_n} - \theta$$

for any character θ of S_n . The result now follows from the branching rule for ordinary representations of S_n (see [7, Ch. 9]). $\square$

This lemma implies that by decomposing the product $\chi^{(n-1,1)}\chi^{(n-1,1)}$ we may find the two rows of X which should be labelled by $\chi^{(n-2,2)}$ and $\chi^{(n-2,1,1)}$. Since these irreducible characters have different degrees (see Lemma 2.2 below) we may fix the row labels $(n - 2, 2)$ and $(n - 2, 1, 1)$.

The remaining two-row characters are found in a similar way. Suppose inductively that there is a unique way to assign the labels $(n - s, s)$ and $(n - s, s - 1, 1)$ for $s \leq r$. If $2r = n$ or $2r + 1 = n$ then we are finished, so we assume that $n > 2r + 1$. By decomposing the product $\chi^{(n-1,1)}\chi^{(n-r,r)}$ we may find the two rows which should be labelled by $\chi^{(n-r-1,r+1)}$ and $\chi^{(n-r-1,r,1)}$. Again, by Lemma 2.2, these characters have different degrees, so we may fix the row labels $(n - r - 1, r + 1)$ and $(n - r - 1, r, 1)$. This completes the inductive step.

Lemma 2.2. *If $r \geq 2$ and $n \geq 2r$ and then*

$$\chi^{(n-r,r)}(1) < \chi^{(n-r,r-1,1)}(1).$$

Proof. By the hook-formula for the irreducible character degrees of S_n (see [6, Theorem 2.3.21]),

$$\chi^{(n-r,r)}(1) = \binom{n}{r} \frac{n-2r+1}{n-r+1}$$

which is less than

$$\chi^{(n-r,r-1,1)}(1) = (r-1) \binom{n}{r} \frac{n-2r+2}{n-r+2}. \quad \square$$

2.3. Fix a column of X . Suppose that in one possible labelling of the columns of X , the corresponding conjugacy class has $g \in S_n$ as a representative. We shall show that the cycle type of g can be reconstructed from the character values $\chi^{(n-r,r)}(g)$, which are known from the previous section. This will show that the column labels of X are uniquely determined.

Since $\chi^{(n-1,1)}(g) = |\text{Fix } g| - 1$, we can find the number of fixed points of g . Suppose inductively that we know that g has a_1 fixed points, a_2 2-cycles, $\dots$, and a_{r-1} $(r-1)$ -cycles, where $1 < r \leq n/2$. Let

$$\pi^{(n-r,r)} = \sum_{s=0}^r \chi^{(n-s,s)}.$$

This is the permutation character of S_n acting on r -subsets of $\{1, \dots, n\}$. Using only the known character values we may calculate $\pi^{(n-r,r)}(g)$, and so find the number of r -subsets fixed by g . This equals the number of r -cycles in g , plus some further quantity that can be computed given $a_1, \dots, a_{r-1}$. Therefore a_r may be determined. Since the action of S_n on r -subsets is isomorphic to its action on $(n-r)$ -subsets, this is sufficient to determine the number of cycles in g of any given length.

2.4. Now we have found all the column labels of X , the remaining row labels are uniquely determined. Here we briefly give a practical way to determine these labels.

It follows from a result of Kramer [9] that two irreducible characters of S_n which agree on every cycle of length r for $1 \leq r \leq n$ are equal. (See [16] for a short proof of this, and some related results.) Suppose then that we have already computed the character values $\chi^\lambda(n-r, 1^r)$ where $0 \leq r < n$ and λ is any partition of n . (One convenient way to do this is to use the Murnagham–Nakayama rule and the hook-formula.) Then to find the partition labelling row i of the character table X , we need only compare the values $X_{i\nu}$ for $\nu = (n-r, 1^r)$ and $0 \leq r < n$ with those in our pre-computed table.

3. ALTERNATING GROUPS

The outer automorphisms of the alternating group A_n induced by the conjugacy action of S_n leads to some inevitable ambiguity in the row and column labels of its character table. Recall that any such automorphism acts as an involution, swapping pairs of split characters and conjugacy classes. More precisely, if λ is a partition of n then $\chi^\lambda \downarrow_{A_n}$ is reducible if and only if λ is self-conjugate, in which case it splits as a sum of two irreducible characters of A_n . Similarly the conjugacy class labelled by the partition ν of n splits in A_n if and only if ν has odd distinct parts. We distinguish split characters

and classes by arbitrarily allocating $+$ and $-$ signs. Whichever allocation we choose, it will be reversed under the outer action of S_n .

For example, the character table of A_5 is

	(1^5)	$(2, 2, 1)$	$(3, 1, 1)$	$(5)^+$	$(5)^-$
(5)	1	1	1	1	1
$(4, 1)$	4	0	1	-1	-1
$(3, 2)$	5	1	-1	0	0
$(3, 1, 1)^+$	3	-1	0	α	β
$(3, 1, 1)^-$	3	-1	0	β	α

where $\alpha = (1 + \sqrt{5})/2$ and $\beta = (1 - \sqrt{5})/2$. The alternative labelling induced by the action of S_5 is given by swapping the characters and conjugacy classes labelled $+$ and $-$.

The next theorem states that this is usually the only alternative labelling.

Theorem 3.1. *Let X be a character table of the alternating group A_n . Provided $n \neq 6$ there is a unique way to assign non-self-conjugate partitions to the rows of X and partitions not having odd distinct parts to the columns of X so that $X_{\lambda\nu} = \chi^\lambda(\nu)$ for all such λ and ν . The labels of the split characters and conjugacy classes are uniquely determined up to signs.*

Proof. For $n \leq 5$ the theorem can be readily verified by inspecting the tables. By [6, Theorem 2.5.15], provided $n \geq 7$, the only character of A_n of degree $n - 1$ is the one labelled by $(n - 1, 1)$. So we can fix this label. Since no self-conjugate partitions appear in the calculations of §2.2, the remaining two row characters may then be identified as before. The cycle types of the columns may also be identified as before. When one comes to a split class the labels $+$ and $-$ may be assigned either way round. Once all the column labels are fixed, there is then a unique way to label the rows. $\square$

Inspection of the character table of A_6 shows that it has 4 alternative labellings. Using the definitions given in §6 below, we have $\text{cAut}(A_6) \cong \chi\text{Aut}(A_6) \cong \text{Out}(A_6) \cong \langle (12), (34) \rangle$.

4. BRAUER CHARACTER TABLES OF SYMMETRIC GROUPS

Let F be a field of prime characteristic p . We say that a partition is p -regular if it has at most $p - 1$ parts of any given size. Recall from [7] that the irreducible FS_n representations are parametrised by the p -regular partitions of n . Let D^μ be the p -modular irreducible corresponding to the p -regular partition μ and let ϕ^μ be the Brauer character of D^μ . (See [11] for an introduction to Brauer characters.) If ν is a partition of n with no part divisible by p then we write $\phi^\mu(\nu)$ for the value of ϕ^μ on the conjugacy class of p' -elements labelled by ν .

In this section we shall prove the following theorem.

Theorem 4.1. *Let X be a Brauer character table of the symmetric group S_n in characteristic p . Unless $n = 6$, or $n = 4$ and $p > 2$, there is a unique way to assign p -regular partitions to the rows of X and partitions with no part divisible by p to the columns of X so that $X_{\mu\nu} = \phi^\mu(\nu)$ for all such μ and ν . In the exceptional cases there are exactly 2 different labellings.*

To prove this theorem we shall need some further results on the modular representations of symmetric groups. Recall that the *decomposition matrix* $D_n(p)$ of S_n in characteristic p is defined by

$$\chi^\lambda = \sum_{\mu} D_n(p)_{\lambda\mu} \phi^\mu$$

when λ is a partition of n and the sum is over all p -regular partitions μ . (Here and elsewhere when we write a relation between ordinary and Brauer characters it is intended to hold for p' -elements only.) Let $\succeq$ denote the dominance order on partitions of n (see [7, Definition 3.2]). The following lemma is Corollary 12.3 in [7].

Lemma 4.2. *Let λ be a partition of n and let μ be a p -regular partition of n . If $D_p(n)_{\lambda\mu} \neq 0$ then $\mu \succeq \lambda$. Moreover $D_p(n)_{\lambda\lambda} = 1$. $\square$*

We also need a simple branching rule for modular representations.

Lemma 4.3. *Let $\mu = (\mu_1, \dots, \mu_k)$ be a partition of n such that $\mu_1 > \mu_2$. If $\bar{\mu} = (\mu_1 - 1, \mu_2, \dots, \mu_k)$ is p -regular then $\phi^\mu \downarrow_{S_{n-1}} = \phi^{\bar{\mu}} + \psi$ where ψ is a sum of Brauer characters ϕ^λ labelled by partitions λ such that $\lambda \triangleright \bar{\mu}$.*

Proof. In future we shall write $\psi \triangleright \mu$ if ψ is a weighted sum of Brauer (or ordinary) characters labelled by partitions λ such that $\lambda \triangleright \mu$. By Lemma 4.2 we may write $\phi^\mu = \chi^\mu - \theta$ where $\theta \triangleright \mu$. It then follows from the ordinary branching rule that

$$\phi^\mu \downarrow_{S_{n-1}} = \chi^{\bar{\mu}} + \psi$$

where $\psi \triangleright \bar{\mu}$. Now apply Lemma 4.2 one more time. $\square$

We can now begin the proof of Theorem 4.1. We follow as closely as possible the method of proof used in §2; thus §4.1 below is the analogue of §2.1, and so on.

4.1. If $n \leq 6$ the theorem may readily be verified by inspecting the tables. (Brauer character tables for $p = 2$ and $p = 3$ and $n \leq 10$ appear in Appendix I.F of [6]; for $p = 5$ and $n = 5, 6$ the required tables may easily be calculated by hand, as all blocks have weight 0 or 1.) The only difference in behaviour from the ordinary case occurs when $n = 4$: the Brauer character table of S_4 in characteristic 2 is shown below.

	(1 ⁴)	(3, 1)
(4)	1	1
(3, 1)	2	−1

Clearly there is no longer any ambiguity about the labels.

4.2. Let $n \geq 7$ and let X be an unlabelled Brauer character table of S_n in characteristic p . As before, it is easy to see that there is a unique way to assign the row label (n) and the column label (1^n) . It was first proved by Wagner (see [13, 14]) that if S is a simple FS_n -module with $\dim S \leq n - 1$ then, provided $n \geq 7$, either S is 1-dimensional, or S is isomorphic to

one of $D^{(n-1,1)}$ or $D^{(n-1,1)} \otimes \text{sgn}$. (For an alternative shorter proof see James [8, Theorem 6].) The Brauer character of $D^{(n-1,1)}$ is

$$\phi^{(n-1,1)}(g) = |\text{Fix } g| - \begin{cases} 1 & \text{if } p \nmid n \\ 2 & \text{if } p \mid n. \end{cases}$$

Of $\phi^{(n-1,1)}$ and $\phi^{(n-1,1)} \text{sgn}$, only $\phi^{(n-1,1)}$ takes $n-4$ as a value, hence we may identify the row of X labelled by $(n-1,1)$.

We are now in a position to identify the rows of X labelled by all two-row partitions. As before, we do this inductively by taking tensor products. However, as there is no simple formula for the degrees of the characters $\phi^{(n-r,r)}$, we have to be slightly more subtle in our approach. The following lemma is the analogue of Lemma 2.1.

Lemma 4.4. *If $r \geq 1$ and $n > 2r + 1$ then*

$$\phi^{(n-1,1)}\phi^{(n-r,r)} = b\phi^{(n-r-1,r+1)} + \phi^{(n-r-1,r,1)} + \psi$$

where $b \geq 1$ and ψ is a weighted sum of irreducible Brauer characters labelled by partitions μ such that $\mu \supseteq (n-r, r-1, 1)$.

Proof. We adapt the argument used in Lemma 2.1. Since $\phi^{(n-1,1)} = 1_{S_{n-1}} \uparrow^{S_n} - c\chi^{(n)}$ for some $c \in \{1, 2\}$, it is sufficient to show that

$$\phi^{(n-r,r)} \downarrow_{S_{n-1}} \uparrow^{S_n} = \phi^{(n-r-1,r+1)} + \phi^{(n-r-1,r,1)} + \psi$$

where $\psi \supseteq (n-r, r-1, 1)$. We have

$$\phi^{(n-r,r)} = \chi^{(n-r,r)} - \theta$$

where $\theta \supseteq (n-r+1, r-1)$. Hence by the ordinary branching rule,

$$\phi^{(n-r,r)} \downarrow_{S_{n-1}} \uparrow^{S_n} = \chi^{(n-r-1,r+1)} + \chi^{(n-r-1,r,1)} + \psi$$

where $\psi \supseteq (n-r, r-1, 1)$. Now apply Lemma 4.2. $\square$

For simplicity we state the following proposition for $p > 2$, and explain later the small modifications needed if $p = 2$.

Proposition 4.5. *Let $n \geq 7$ and let C be a labelled Brauer character table of S_n in characteristic $p > 2$. Suppose that the first column is labelled by (1^n) , and that the rows are arranged so that the first row is labelled by (n) , the second by $(n-1, 1)$ and the next $2(r-1)$ with the labels*

$$(n-2, 2), (n-3, 3), \dots, (n-r, r) \\ (n-2, 1^2), (n-3, 2, 1), \dots, (n-r, r-1, 1).$$

in any order. If we are given the first $2r$ rows of C with the row and column labels removed, then the row labels may be uniquely reconstructed.

Proof. We work by induction on n . If $n = 7$ and $p = 3$ or $p = 5$ then the Brauer characters of S_n that can appear in X have distinct degrees, so the result is immediate. If $p = 7$ then both $\phi^{(5,2)}$ and $\phi^{(4,3)}$ have degree 14, but only the former takes 6 as a value. (As both characters are in blocks of weight zero this can be seen directly from the ordinary character table.)

Suppose now that $n \geq 8$. By hypothesis, we may immediately attach the row labels (n) and $(n-1, 1)$ to C . We now attempt to reach a situation in

which the inductive hypothesis for $n-1$ can be applied. Notice first that the values of $\phi^{(n-1,1)}$ determine which columns in the table come from conjugacy classes with at least one fixed point, and so are relevant when we restrict a character to S_{n-1} . The restriction of $\phi^{(n)}$ to S_{n-1} is of course $\phi^{(n-1)}$. We may obtain $\phi^{(n-2,1)}$ by removing any copies of $\phi^{(n-1)}$ from $\phi^{(n-1,1)} \downarrow_{S_{n-1}}$.

By Lemma 4.4 above, when we express $\phi^{(n-1,1)}\phi^{(n-1,1)}$ as a sum of rows of X , two new characters appear: $\phi^{(n-2,2)}$ and $\phi^{(n-1,1,1)}$. When we restrict these new characters to S_{n-1} we get, in addition to any copies of $\phi^{(n-2,1)}$ and $\phi^{(n-2)}$ that may be present, two new Brauer characters of S_{n-1} : namely $\phi^{(n-3,1,1)}$ and $\phi^{(n-3,2)}$. We may now apply the inductive hypothesis (with $r = 2$) to determine which label should go with which. To get back to S_n we use Lemma 4.3. Together with Lemma 4.2, it implies that $\phi^{(n-2,2)} \downarrow_{S_{n-1}}$ does not contain $\phi^{(n-3,1,1)}$, whereas $\phi^{(n-1,1,1)} \downarrow_{S_{n-1}}$ does. We use this to fix the labels $(n-2, 2)$ and $(n-2, 1, 1)$.

The remaining two-row labels are fixed by repeating this argument, in a way closely analogous to the proof of Lemma 2.2. We therefore leave the remaining details of the proof to the reader. $\square$

By a further induction on r we may use this proposition to identify the rows of X labelled by two-row partitions.

If $p = 2$ then the statement of Proposition 4.5 must be slightly modified. If n is odd we must delete $(n-2, 1, 1)$, and if $n = 2m$ is even we must delete $(n-2, 1, 1)$ and (m, m) , as these partitions are no longer p -regular. The main change in the proof is that now $\phi^{(n-1,1)}\phi^{(n-1,1)}$ only contains one new Brauer character, $\phi^{(n-2,2)}$; this makes the first step is slightly simpler. After that, no alterations are needed, unless $n = 2m$ is even, in which case the last two row Brauer character we must find is $\phi^{(m+1, m-1)}$. Again if anything this makes the process slightly simpler.

4.3. We now determine the column labels of X . If we order the rows and columns of the decomposition matrix $D_n(p)$ by the dominance order, (n) appears first of all, followed by the two-row partitions of n . By Lemma 4.2, $D_n(p)_{\lambda\mu} = 0$ if λ has at most two rows and μ does not. Hence the matrix $D_n(p)X$ has at its top the values of the characters χ^λ for partitions λ with at most two rows.

We can now use the same argument as in the ordinary case to determine the column labels of X . Once we have a complete set of column labels the row labels are, of course, fixed. This completes the proof of Theorem 4.1.

5. BRAUER CHARACTER TABLES OF ALTERNATING GROUPS

Provided p is odd, the modifications to the work of §4 needed to deal with the Brauer character tables of alternating groups in characteristic p are analogous to those needed to the work of §2 (and made in §3) to deal with the ordinary character tables of alternating groups.

Let X be a Brauer character table of the alternating group A_n in odd-characteristic. As in §3, we label the split characters and conjugacy classes by $+$ and $-$ signs. As usual, to get started we need to identify the character labelled by $(n-1, 1)$. For this we use Theorem 1.1 in [14], which states that if $n \geq 7$ and ϕ is an odd-characteristic Brauer character of A_n such

that $\phi(1) \leq n$ then $\phi = \phi^{(n-1,1)} \downarrow_{A_n}$. (For an alternative shorter proof for $n \geq 10$ see [8, Theorem 7(ii)]; the result can easily be checked directly in the remaining cases.)

We also need to know that none of the characters considered in §4.2 split on restriction to A_n . For this to hold we must take $n \geq 8$.

Lemma 5.1. *If λ is a p -regular partition of n let $m(\lambda)$ be the p -regular partition of n defined by*

$$D^\lambda \otimes \text{sgn} = D^{m(\lambda)}.$$

If either λ has at most two rows, or λ is of the form $(n-r, r-1, 1)$ for some $n \geq 8$ and $r \leq n/2$ then $\lambda \neq m(\lambda)$. Hence ϕ^λ does not split on restriction to A_n .

Proof. It follows easily from Ford's description of the Mullineux map in [4] that $\lambda \neq m(\lambda)$ if either of the conditions on λ hold. Ford's paper also gives the Clifford theory needed to prove the second assertion. $\square$

It is not possible to take $n \geq 7$ (as was the case in §4.2) because if $p = 3$ then $m((4, 2, 1)) = (4, 2, 1)$. The base case in the analogue of Proposition 4.5 is therefore $n = 8$. Calculation shows that the Brauer characters of A_8 that can appear in the table C have distinct degrees when $p = 3$ and when $p = 7$. If $p = 5$ then $\phi^{(6,1,1)} = \phi^{(5,3)} = 21$, but only the former character takes 6 as a value, so again there is no ambiguity. Thus we may identify the rows of X labelled by two-row partitions. The column labels may now be determined in essentially the same way as §4.3.

Direct examination of the cases for $n \leq 7$ gives the following theorem.

Theorem 5.2. *Let X be a Brauer character table of the alternating group A_n in odd characteristic p . Provided $n \neq 6$ there is a unique way to assign p -regular partitions λ such that $m(\lambda) \neq \lambda$ to the rows of X and partitions not all of whose parts are odd, and with no part divisible by p , to the columns of X so that $X_{\lambda\mu} = \phi^\lambda(\mu)$ for all such λ and μ . The labels of the split characters and conjugacy classes are uniquely determined up to signs.* $\square$

When $n = 6$ and $p = 3$ there are two different labellings, interchanged by the conjugacy action of S_6 . When $n = 6$ and $p = 5$ there are again two different labellings, but this time S_6 acts trivially, and they are interchanged only by the outer automorphism of S_6 .

6. A MORE GENERAL SETTING

Given an arbitrary finite group G , there is usually no canonical way to label the rows and columns of its character table. However, as the reader familiar with Brauer's permutation lemma will already have realised, this need not stop us from considering analogous versions of our results.

Let k be the number of conjugacy classes of G , and let X be a character table of G . We say that a pair $(\sigma, \tau) \in S_k \times S_k$ is an *automorphism* of X if $X_{i\sigma j\tau} = X_{ij}$ whenever $1 \leq i, j \leq k$. Let $\text{Aut}(X)$ be the group of all automorphisms of X . It is clear that for each $\sigma \in S_k$ there is at most one $\tau \in S_k$ such that $(\sigma, \tau) \in \text{Aut}(X)$. We may therefore define a group $\chi\text{Aut}(G)$ by

$$\chi\text{Aut}(G) = \{\sigma \in S_k : (\sigma, \tau) \in \text{Aut}(X) \text{ for some } \tau \in S_k\}.$$

This group is well-defined up to conjugacy in S_k . For example, our Theorem 1.1 states that $\chi\text{Aut}(S_n)$ is trivial unless $n = 4$ or $n = 6$.

Problem 6.1. *Calculate $\chi\text{Aut}(G)$ for important classes of groups.*

In connection with this problem, it is interesting to explore the relationship between $\chi\text{Aut}(G)$ and $\text{Out}(G) = \text{Aut}(G)/\text{Inn}(G)$, the group of outer automorphisms of G . Clearly there is a group homomorphism

$$c : \text{Out}(G) \rightarrow \chi\text{Aut}(G)$$

defined for $\gamma \in \text{Out}(G)$ by letting $c(\gamma)$ be the permutation induced by γ on the ordinary characters of G . In some cases c is an isomorphism — for example, this is the case if G is abelian, or G is a symmetric group other than S_4 , or G is any alternating group. But, as the example of S_4 shows, c need not be surjective. The dual question, of whether c must be injective, or equivalently, whether there is a finite group G and an outer automorphism $\gamma \in \text{Aut}(G)$ such that $\chi^\gamma = \chi$ for all irreducible characters χ , was considered by Burnside: see Note B in [3]. In [15], G. E. Wall gives an example in which G has order 32 and γ has order 4.

It is worth noting that Burnside's question can be stated without even mentioning characters, since by Brauer's permutation lemma (see [2, §6]), t is such an automorphism if and only if t permutes within themselves all the conjugacy classes of G . (Incidentally, it seems clear from [3, §186] that Brauer's permutation lemma was already well known to Burnside.)

Another obvious question, which is related to Problem 6.1, is:

Problem 6.2. *Is the map $c : \text{Out}(G) \rightarrow \chi\text{Aut}(G)$ always an isomorphism when G is a finite simple group?*

A final problem, which can be answered more easily, arises from the definition of $\chi\text{Aut}(G)$. If we look instead at the admissible *column* permutations of X then we obtain the group

$$\text{cAut}(G) = \{\tau \in S_k : (\sigma, \tau) \in \text{Aut}(X) \text{ for some } \sigma \in S_k\}.$$

By Brauer's permutation lemma the groups $\chi\text{Aut}(G), \text{cAut}(G) \leq S_k$ are isomorphic as abstract groups, *via* an isomorphism preserving the cycle types of elements. But this on its own does not guarantee that they are permutation isomorphic, as the two subgroups of S_6 ,

$$\langle (12)(34), (13)(24) \rangle, \quad \langle (12)(34), (12)(56) \rangle$$

show. (There are many more examples of this type.) The following example shows that $\chi\text{Aut}(G)$ and $\text{cAut}(G)$ need *not* be permutation isomorphic, and so we made a genuine choice in concentrating on χAut earlier.

Example 6.3. Let $G \cong C_2 \times D_8$, where D_8 is the dihedral group of order 8. The character table of G is, with one ordering of the rows and columns:

-1	1	1	-1	1	-1	-1	-1	1	1
-1	-1	1	1	-1	1	-1	-1	1	1
1	-1	-1	1	1	-1	-1	-1	1	1
1	1	-1	-1	-1	1	-1	-1	1	1
-1	1	-1	1	-1	-1	1	1	1	1
1	-1	1	-1	-1	-1	1	1	1	1
-1	-1	-1	-1	1	1	1	1	1	1
0	0	0	0	0	0	-2	2	2	-2
0	0	0	0	0	0	2	-2	2	-2
1	1	1	1	1	1	1	1	1	1

One finds that

$$\begin{aligned}\chi\text{Aut}(G) &= \langle (1234)(56), (34)(57), (89) \rangle, \\ \text{cAut}(G) &= \langle (1234)(56), (25)(46), (78) \rangle\end{aligned}$$

where the isomorphism $\chi\text{Aut}(G) \cong \text{cAut}(G)$ is indicated by the order of generators. As the orbits of $\chi\text{Aut}(G)$ have sizes 4, 3, 2, 1 whereas the orbits of $\text{cAut}(G)$ have sizes 6, 2, 1, 1, the two groups are not permutation isomorphic. (Abstractly, both are isomorphic to $S_4 \times C_2$.)

Finally we mention a theorem of Higman (see [5, Theorem 8.21]) which states that given a character table of a finite group, one can determine prime divisors of the orders of the group elements corresponding to any given column. It is well known that the dihedral and quaternion groups of order 8 have the same character table, so this is the most one can hope for in general. Theorem 1.1 and Theorem 3.1 imply that for symmetric and alternating groups much more is true.

Corollary 6.4. *Given an unlabelled character table of a symmetric group other than S_4 one may determine the order of the elements corresponding to any of its columns. The same result holds for any alternating group.* $\square$

ACKNOWLEDGEMENT

I should like to thank Dr Christine Bessenrodt for the helpful comments she made when this work was beginning.

REFERENCES

- [1] BESSENRODT, C., AND UNO, K. Character relations and simple modules in the Auslander-Reiten graph of the symmetric and alternating groups and their covering groups. *Algebr. Represent. Theory* 4, 5 (2001), 445–468.
- [2] BRAUER, R. On the connection between the ordinary and the modular characters of groups of finite order. *Ann. of Math. (2)* 42 (1941), 926–935.
- [3] BURNSIDE, W. *Theory of groups of finite order*. Dover Publications Inc., New York, 1955. 2nd ed.
- [4] FORD, B. Irreducible representations of the alternating group in odd characteristic. *Proc. Amer. Math. Soc.* 125, 2 (1997), 375–380.
- [5] ISAACS, I. M. *Character theory of finite groups*. Dover Publications Inc., New York, 1994. Corrected reprint of the 1976 original [Academic Press, New York; MR0460423 (57 #417)].

- [6] JAMES, G., AND KERBER, A. *The representation theory of the symmetric group*, vol. 16 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley Publishing Co., Reading, Mass., 1981.
- [7] JAMES, G. D. *The representation theory of the symmetric groups*, vol. 682 of *Lecture Notes in Mathematics*. Springer, Berlin, 1978.
- [8] JAMES, G. D. On the minimal dimensions of irreducible representations of symmetric groups. *Math. Proc. Cambridge Philos. Soc.* 94, 3 (1983), 417–424.
- [9] KRAMER, P. Factorisation of projection operators for the symmetric group. *Z. Naturforsch 21A* (1966), 657–658.
- [10] NAGAO, H. On the groups with the same table of characters as symmetric groups. *J. Inst. Polytech. Osaka City Univ. Ser. A.* 8 (1957), 1–8.
- [11] NAVARRO, G. *Characters and blocks of finite groups*, vol. 250 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1998.
- [12] ROTMAN, J. J. *An introduction to the theory of groups*, fourth ed., vol. 148 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [13] WAGNER, A. The faithful linear representation of least degree of S_n and A_n over a field of characteristic 2.. *Math. Z.* 151, 2 (1976), 127–137.
- [14] WAGNER, A. The faithful linear representations of least degree of S_n and A_n over a field of odd characteristic. *Math. Z.* 154, 2 (1977), 103–114.
- [15] WALL, G. E. Finite groups with class-preserving outer automorphisms. *J. London Math. Soc.* 22 (1947), 315–320 (1948).
- [16] WILDON, M. Character values and decomposition matrices of symmetric groups. *Submitted. Available at ArXiv math.RT/0610414* (2006).

E-mail address: m.j.wildon@swansea.ac.uk